INDICATION Predo for the symptomatic treatment of

inflammatory processes. For the symptomatic treatment of allergic read and anti-toxic effects. It is primarily indicated for arthritis, synovitis, inflammation of tendon shea strains, itching, eczema, dermatitis, and as an

adjuvant in infectious diseases. STORAGE: Store in a cool place protected from light and below 25 C.

For Veterinary use Only

\*\*Keep out of reach of children



SUSPENSION INJECTABLE IV. IM

Prednisolon Excipient

PHARMACEUTICAL FORM

# DOSAGE

horses: 10ml / animal Dogs: 0.5 - 1 ml / animal







## PREDO 0.5

### COMPOSITION (Per ml):

Prednisolone Acetate: 5 mg Excipients: q.s. to 1 ml.

#### PHARMACEUTICAL FORM:

Sterile injectable suspension for intravenous (IV) or intramuscular (IM) administration.

#### INDICATIONS:

Predo 0.5 is indicated for use in horses and dogs for the management of:

- Musculoskeletal inflammation (arthritis, tendinitis, bursitis)
- Allergic conditions (dermatitis, urticaria, respiratory allergies)
- · Shock and stress-related reactions
- · Autoimmune and inflammatory diseases
- Supportive treatment in acute infections (alongside antibiotics).

#### DOSAGE AND ADMINISTRATION

 Species
 Dosage Range
 Route

 Horses:
 0.05-0.1 mg/kg body weight
 IV or IM

 Dogs:
 0.25-0.5 mg/kg body weight
 IM

Shake well before use. Use sterile equipment only.

#### PRECAUTIONS:

- Avoid use in animals with systemic fungal or viral infections.
- Do not mix with other injections in the same syringe.
- Prolonged or repeated use may cause adrenal suppression or metabolic effects.
- · Use only under veterinary supervision.

#### STORAGE:

Store below 25°C. Protect from light and freezing. Keep out of reach of children.

#### PRESENTATION:

Multidose vial - 50 ml.

### MANUFACTURER:

Credence Remedies Pvt. Ltd.
Office No. 313, 3rd Floor, Puranik Capitol,
Behind Vijay Park, Kasarvadavali, Ghodbunder Rd,
Thane (W) – 400615, Maharashtra, India.